

# Complementary Food Hygiene

An overlooked opportunity in the WASH, nutrition and health sectors

POLICY BRIEF



Mothers publicly committing to practise food hygiene behaviours in the Nepal intervention-study. Credit: LSHTM/Om Prasad Gautam

## Summary

Poor complementary food hygiene may account for a substantial proportion of diarrhoeal diseases among infants and young children in developing countries, which in turn are responsible for a large share of the global burden of infectious disease. However, most of the information on food hygiene in developing countries derives from expert opinion and biological plausibility rather than robust epidemiological evidence. This briefing paper documents the Sanitation and Hygiene Applied Research for Equity (SHARE) consortium's contribution to narrowing the evidence gap, highlights opportunities for future research, and offers insights that could influence policy and improve programming in the water, sanitation and hygiene (WASH), nutrition and health sectors globally.

### SHARE Consortium

The SHARE consortium is a five year initiative (2010-2015), funded by the UK Department for International Development, which seeks to contribute to accelerated progress on the Millennium Development Goal sanitation target by generating critical knowledge to inform policy and practice in the areas of sanitation and hygiene.

**Written by:**  
Om Prasad Gautam<sup>1</sup>  
Joanna EstevesMills<sup>1</sup>  
Alexandra Chitty<sup>1</sup>  
Val Curtis<sup>1</sup>

<sup>1</sup>London School of Hygiene & Tropical Medicine

## Complementary foods: a key disease transmission pathway

Diarrhoeal diseases are the second leading cause of death for children under the age of five globally [1], and kill more young children than AIDS, malaria and measles combined [2]. They also increase the risk of malnutrition [3, 4], which itself accounts for almost a quarter of the global disease burden for children under the age of five. It should be noted, however, that diarrhoeal diseases are not the only factor in the causal pathway between faecal contamination and undernutrition; malabsorption, caused by environmental enteropathy and nematode infections, also plays a role [5].

Contaminated complementary foods may account for a substantial proportion of diarrhoeal diseases among infants and young children in developing countries [6]. While it is important for an infant's development to supplement breast milk with appropriate solid foods from six months, unhygienic preparation and feeding of complementary foods exposes infants in low income settings to pathogens of faecal origin. There is evidence that the incidence of diarrhoeal disease is higher in children after weaning is initiated [7]. In fact, in low income settings, the level of contamination in complementary foods can be higher than in drinking water [8], though this varies between environmental settings [9-11]. Furthermore, children under the age of six months may also be exposed to such pathogens as a result of non-exclusive breastfeeding [12].




Figure 1: 'F' diagram illustrating transmission pathways.

*Credit: Om Prasad Gautam*

Interventions to improve public health have not adequately addressed the food contamination pathway. The nutrition sector has traditionally placed greater emphasis on 'nutrition specific' than on 'nutrition sensitive'<sup>1</sup>[13] interventions, and therefore the evidence base on the impact of the latter on tackling the underlying determinants of foetal and child nutrition and development is limited. Meanwhile, although hygiene is high on the agenda in the WASH sector, interventions have often been limited to hand washing

<sup>1</sup> 'Nutrition-specific' programmes address the immediate determinants of foetal and child nutrition and development, whilst 'nutrition-sensitive' programmes address the underlying determinants.

There are several reasons for this neglect:

- Health institutions tend to prioritise vertical disease management programmes over horizontally integrated public health efforts such as prevention programmes;
- The nutrition sector places greater emphasis on dietary intake than on food hygiene;
- There is a lack of evidence on the effect of food hygiene interventions on child health outcomes; Guidance on best practice for securing sustainable food hygiene behaviour change in various settings is lacking.

## The evidence base

There is evidence to suggest that unhygienic food preparation and feeding is likely to increase the level of microbiological contamination in food and the risk of diarrhoeal disease [6, 8, 14, 15]. There is also some evidence identifying risk factors and associated behaviours. However, little is known about food hygiene practices in low income settings or about interventions that could mitigate the risks [16].

### What We Know

Studies to date have focused on the factors that may lead to food-borne contamination. These include: hot climate [8], poor storage practices, insufficient cooking time [6, 8], time elapsed between meal preparation and feeding [17, 18], and use of unsterilized and dirty feeding bottles for children [7, 8].

Expert opinion, meanwhile, suggests that risk factors for food-borne contamination include: environmental contamination due to lack of sanitation; use of contaminated water to wash serving utensils; not washing hands prior to cooking and feeding; consumption of food that has been spilled on the floor; and use of dirty cloths for wiping hands and utensils.

The hygiene practices of mothers have been found to be related to a high level of bacterial contamination of drinking water and complementary foods [19], and the home has been identified as an important location for acquiring food-borne diseases [20], due, in large part, to specific food hygiene practices [17].

We also know that improving socially or culturally engrained practices is challenging and requires going beyond purely supply-led interventions [21]. Evidence suggests that addressing the structural determinants of hygiene practices will be integral to securing sustainable food hygiene behaviour change [22, 23].

When SHARE began in 2010, effective food hygiene interventions had not been described or evaluated, though the Hazard Analysis and Critical Control Point (HACCP) approach had been used to identify points where controls could be applied to prevent, reduce or eliminate this contamination [24-27]. One study in peri-urban Bamako, Mali, had gone further and had designed

and delivered a small-scale hygiene experiment using HACCP which proved to be effective in reducing the contamination of complementary foods [24].

### The evidence gap

In 2010, many aspects of the relationship between food hygiene and child health remained unclear and there was limited knowledge about how best to design and deliver interventions. Firstly, our assumptions about food-borne infections in low income settings were still largely based on expert opinion and biological plausibility, rather than data from epidemiological studies [16]. There was a need to better characterise pathogen transmission through food and to establish how this differed across settings. Secondly, it remained unclear which factors, or behaviours, posed the greatest risk of food contamination and should therefore be targeted by interventions seeking to reduce these risks. SHARE was also keen to establish how to promote such interventions in a normal community programme setting, and to identify how best to tackle multiple behaviour change.

## Narrowing the evidence gap

Recognising the importance of food hygiene in the prevention and control of faecally transmitted diseases and the need to build a stronger evidence base, SHARE provided funding for three studies in Bangladesh, Nepal and the Gambia.

### Bangladesh

Using the same principles employed in the Mali study and based on the results [24], SHARE funded a small intervention study to investigate whether hygiene promotion following the HACCP approach was effective in reducing complementary food contamination in Bangladesh. In the study a total of 60 households in rural Matlab, Bangladesh, were selected (30 each in the control and treatment arms). Two types of complementary foods were collected from each household and analysed for microbial contamination just after cooking and before the child was fed. Critical control points were determined, and mothers in the study households were trained for four weeks to achieve and monitor these. Results showed that the hygiene intervention significantly reduced food contamination. The study highlighted the importance of complementary foods as an exposure pathway for enteric pathogens [9]. It also demonstrated that controlling critical points of transmission can significantly reduce this exposure.

## Nepal

Building on the Mali and Bangladesh findings, SHARE funded the design, delivery and evaluation of an intervention to change the food hygiene behaviours of mothers in rural Nepal [28]. This consisted of a motivational package targeting five key food hygiene behaviours using emotional drivers rather than cognitive appeals, as well as emphasis on change in behaviour settings. Four clusters received the intervention over a period of three months whilst four clusters remained a control group. Outcomes were measured in 239 households with a child aged 6-59 months. 43% of mothers were able to maintain all 5 key behaviours and the intervention was successful in significantly improving the microbiological indicators in complementary foods. The results suggest that it is possible to substantially improve food hygiene behaviour and reduce the risk of microbiological contamination of complementary foods through scalable community level interventions.



Five key food hygiene behaviours promoted in Nepal.  
Credit: Om Prasad Gautam

## The Gambia

An ongoing SHARE-funded cluster randomised control trial in rural Gambia build on the lessons learned from earlier SHARE-funded work on food hygiene and behaviour change, including the successful SuperAmma campaign in India [29]. It aims to substantiate the findings of the Nepal study through adaptation of the intervention to a different context. As in Nepal, it seeks to provide an intervention that can be scaled-up, this time in low income settings in Africa<sup>1</sup>. Formative research has been completed and the intervention is in the process of being refined.

<sup>1</sup> The SuperAmma trial was a communication campaign based on the Evo-Eco theory of behaviour change that was carried out in Southern India to inculcate the habit of handwashing with soap. Further information about SuperAmma can be found here: <http://www.superamma.org/index.html>.



This package of research has plugged an important evidence gap in the sector. The studies in Mali and Bangladesh showed that food contamination can be reduced using a HACCP methodology at the household level [9]. The study in Nepal, meanwhile, successfully tested a simple and scalable food hygiene intervention targeting multiple behaviours that could be implemented in a community setting through health, WASH and nutrition programmes [28]. We await results from the Gambia study with anticipation; these should be available in early 2015.

## More knowledge is still needed

These contributions to our knowledge of this important area for child health are already contributing to changes in policy and practice. In Nepal, for example, a pilot project is exploring how hygiene, including food hygiene, can be integrated into vaccination campaigns [30]. However, major knowledge gaps remain:

- **Changing behaviour:** Getting mothers to change long-held routine kitchen practices will always be difficult. We need clever new ways, informed by socio-ecological approaches to behaviour change - such as those piloted in Nepal - to secure sustainable food hygiene behaviour change.
- **Transmission pathways:** We need to fully understand each of the transmission pathways of all of the major enteric pathogens, including through complementary foods, in different settings so as to be able to better target our interventions.
- **Child health:** The effect of food-borne contamination on child health during the critical window of weaning and long term development needs to be better understood. In particular, the relationship between food-borne pathogens and malnutrition needs more evidence. This would inform policy at national and international levels.
- **Scaling-up:** Further development and testing of strategies for scaling-up complementary food hygiene interventions - such as that trialled in Nepal - are required. This would include working closely with nutrition programmes already targeting improvements in complementary feeding practices at scale and with WASH programmes where hygiene components have been included but have not specifically focussed on food hygiene. Scaling-up interventions, and the cross-sectoral collaboration associated therein, would lead to better outcomes for children and would increase the benefits from investment in nutrition, health and WASH.

## Recommendations

Whilst our knowledge remains incomplete, the current state of evidence suggests that it is worthwhile investing in food hygiene and, more specifically, in carefully designed programmes to improve food hygiene behaviour. The successful improvement of safe food hygiene practices can help to reduce or eliminate complementary food contamination, resulting in the prevention or control of diarrhoea and malnutrition. The following recommendations would likely have a major impact in ensuring this:

1

Food hygiene interventions must be carefully designed based on scientific principles of behaviour change.

Evidence from other sectors suggests that simply lecturing and messaging is not effective [22] whilst programmes that creatively target emotions and the settings in which behaviour takes place are likely to be more successful [28, 29].

2

There must be better integration across the nutrition, health and WASH sectors nationally and internationally, and food hygiene should be mainstreamed.

Evidence-based international guidelines that better reflect the complex determinants of food hygiene behaviours and suggest greater cross-sectoral collaboration, are essential. Ongoing post-2015 discussions provide the perfect setting and crucial range of stakeholders required for such efforts. Guidelines formulated could then be adapted to the national context through a participatory process that engages all relevant stakeholders.

3

High quality research in this area must continue to be funded and carried out.

Additional large-scale research is needed to better understand this transmission pathway, how it affects child health and long term development, and to further test scalable behaviour change interventions in different settings.

## Recommendations

**4**

There needs to be better, more consistent monitoring of complementary food contamination and its causes across the sectors concerned.

Continual monitoring and surveillance of food-borne infection, its outbreaks, and associated causal factors is lacking in low income settings, but is essential for ensuring adequate quantification and understanding of the problem. Joint-sectoral indicators must be established and food hygiene components should be included in Demographic Health Surveys, national level nutrition plans, health sector information management systems, and ongoing WASH monitoring mechanisms. Furthermore, disease surveillance mechanisms should disaggregate according to the different pathogens that can cause food-borne infection.

**5**

**Governments, donors and practitioners must increase their financing of complementary food hygiene interventions.**

The damage caused by undernutrition to a child before the age of two is irreversible [31] and goes on to predict a range of health and other outcomes later in life [32, 33]. Therefore, increased investment in complementary food hygiene is critical to child health and development.

**6**

**Financial increases must be accompanied by better targeting and greater efforts to ensure financial absorption.**

In low income settings, the targeting of often scarce resources is frequently inadequate in the nutrition and WASH sectors. Greater attention must be paid to, and targeting should reflect, the disease epidemiology and transmission pathways of food-borne infections. Furthermore, financial increases should go hand in hand with improved budget execution and relevant human resource capacity building.



## References

1. Liu, L., et al., Global, regional, and national causes of child mortality: an updated systematic analysis for 2010 with time trends since 2000. *Lancet*, October 2012 379(9832): p. 2151-2161.
2. UNICEF/WHO, Diarrhoea: why children are still dying and what can be done. 2009.
3. Briend, A., Is diarrhoea a major cause of malnutrition among the under-fives in developing countries? A review of available evidence. *Eur J Clin Nutr* 44(9):611-28., 1990.
4. Checkley, W., et al., Multi-country analysis of the effects of diarrhoea on childhood stunting. *International Journal of Epidemiology*, 2008. 37: p. 816-830.
5. O'Lorcain, P. and C. Holland, The public health importance of *Ascaris lumbricoides*. *Parasitology*, 2000: p. S51-71.
6. Motarjemi, Y., et al., Contaminated weaning food: a major risk factor for diarrhoea and associated malnutrition. *Bulletin of the World Health Organization* 1993. 71 (1): p. 79-92.
7. Barrel, R. and M. Rowland, Infant foods as a potential source of diarrhoeal illness in rural West Africa. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 1997(73): p. 85-89.
8. CF. Lanata, Studies of food hygiene and diarrhoeal disease. *International Journal of Environmental Health Research*, 2003. 13:S1: p. S175-S183.
9. Islam, M., et al., Hygiene intervention reduces contamination of weaning food in Bangladesh. *Tropical Medicine and International Health*, March 2013. volume 18 no 3: p. pp 250-258.
10. Toure, O., et al., Improving microbiological food safety in peri-urban Mali; an experimental study. *Food Control*, 2011. 22: 10 p. 1565-1572. 25 ref.
11. Ngure, F., et al., Formative Research on Hygiene Behaviors and Geophagy among Infants and Young Children and Implications of Exposure to Fecal Bacteria. *American Journal of Tropical Medicine and Hygiene*, 2013. 89(4): p. 709-716.
12. Cai, X., T. Wardlaw, and D. Brown, Global trends in exclusive breastfeeding. *International Breastfeeding Journal*, 2012. 7(12).
13. Ruel, M., H. Alderman, and Maternal and Child Nutrition Study Group, Nutrition-sensitive interventions and programmes: how can they help to accelerate progress in improving maternal and child nutrition? *Lancet* 2013; 382: , 2013. 382: p. 536-51.
14. Potgieter, N., et al., Bacterial contamination of Vhuswa--a local weaning food and stored drinking-water in impoverished households in the Venda region of South Africa. *Journal of Health, Population and Nutrition*, 2005. 23: 2: p. 150-155.
15. Sheth, M. and R. Dwivedi, Complementary foods associated diarrhea. *Indian Journal of Pediatrics*, 2006. 73(1): p. 61-4.

16. Curtis, V., et al., Hygiene: new hopes and new horizons. *Lancet Infect Dis.*, 2011. 11: p. 312-21.
17. Black, R., et al., Incidence and etiology of infantile diarrhoea and major routes of transmission in Huascar, Peru. *American Journal of Epidemiology*, 1989. 129 (4): p. 785-799.
18. Molbak, K., et al., Bacterial contamination of stored water and stored food: a potential source of diarrhoeal disease in West Africa. *Epidemiology & Infection*, 1989 Apr 102(2): p. 309-16.
19. Imong, S., et al., The bacterial content of infant weaning foods and water in rural northern Thailand. *Journal of Tropical Pediatrics*, 1989. 35(1): p. 14-8.
20. Redmond, E. and C. Griffith, The importance of hygiene in the domestic kitchen: implications for preparation and storage of food and infant formula. *Perspectives in Public Health*, 2009 March. 129(2): p. 69-76.
21. Jenkins, M. and S. Sugden, Rethinking Sanitation: Lessons and Innovation for Sustainability and Success in the New Millennium. *Human Development Report 2006*. Human Development Report Office Occasional Paper. , 2006.
22. Biran, A., et al., The effect of a soap promotion and hygiene education campaign on handwashing behaviour in rural India: a cluster randomised trial. *Tropical Medicine and International Health* 2009. 14(10): p. 1303-1314.
23. Curtis, V., et al., Planned, motivated and habitual hygiene behaviour: an eleven country review. *Health Education Research Advance Access*, March, 2009.
24. Toure, O., et al., Piloting an intervention to improve microbiological food safety in Peri-Urban Mali. *Int. J. Hyg. Environ. Health* (2012), 2013. doi:10.1016/j.ijheh.2012.02.003.
25. Sheth, M., et al., Hazard analysis and critical control points of weaning foods. *Indian Journal of Pediatrics*, 2000 Jun. 67(6): p. 405-10.
26. Oranusia, S., V. Umoha, and J. Kwagab, Hazards and critical control points of kunun-zaki, a non-alcoholic beverage in Northern Nigeria. *Food Microbiology*, 2003. 20: p. 127-132.
27. Motarjemi, Y. and M. Schothorst, HACCP Hazard Analysis and Critical Control Point Principles and Practice. *Teacher's Handbook*. World Health Organization, 1999. WHO/SDE/PHE/FOS/99.3.
28. Gautam, O., et al., Effect of an intervention to improve five food hygiene behaviours of mothers in Nepal: an exploratory trial. (in process of submission - not yet published), 2014.
29. Biran, A., et al., Effect of a behaviour-change intervention on handwashing with soap in India (SuperAmma): a cluster-randomised trial. *The Lancet Global Health*, 2014. 2(3): p. e145-e154.

30. Velleman, Y., K. Greenland, and O. Gautam, An opportunity not to be missed - immunization as an entry point for hygiene promotion and diarrhoeal diseases reduction in Nepal. *Journal of Water, Sanitation and Hygiene for Development* 2013. 1(03.3): p. 459-466.
31. Martorell, R., L. Khan, and D. Schroeder, Reversibility of stunting: epidemiological findings in children from developing countries. *Eur J Clin Nutr*, 1994: p. 1:S45-57.
32. Victora, C., et al., Maternal and child undernutrition: consequences for adult health and human capital. *Lancet*, 2008. 371(9609): p. 340-357.
33. Horton, R., Maternal and child undernutrition: an urgent opportunity. *Lancet*, 2008. 371(9608): p. 179.

## Research for sanitation and hygiene solutions

The world is seriously off-track in meeting the Millennium Development Goal on sanitation and 2.6 billion people are still without a safe toilet.

SHARE aims to address these challenges by accelerating progress on sanitation and hygiene in developing countries by generating rigorous and relevant research and ensuring new and existing solutions are adopted at scale.

The consortium conducts research across four pillars:

- Health
- Equity
- Urban
- Markets

SHARE's activities primarily take place in its focus countries:

- Bangladesh
- India
- Malawi
- Tanzania

The DFID-funded SHARE consortium is led by the London School of Hygiene & Tropical Medicine. Its other partners are the International Centre for Diarrhoeal Disease Research, Bangladesh, International Institute for Environment and Development, Shack/Slum Dwellers International and WaterAid.

January 2015



This material has been funded by UK aid from the Department for International Development (DFID). However, the views expressed do not necessarily reflect the Department's official policies.

---

### Contributor

LONDON  
SCHOOL of  
HYGIENE  
& TROPICAL  
MEDICINE

